



Relation between *in Utero* Arsenic Exposure and Birth Outcomes in a Cohort of Mothers and Their Newborns from New Hampshire

Diane Gilbert-Diamond, Jennifer A. Emond, Emily R. Baker,
Susan A. Korrick, and Margaret R. Karagas

<http://dx.doi.org/10.1289/ehp.1510065>

Received: 10 April 2015

Accepted: 12 February 2016

Advance Publication: 8 March 2016

Note to readers with disabilities: *EHP* will provide a [508-conformant](#) version of this article upon final publication. If you require a 508-conformant version before then, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.



National Institute of
Environmental Health Sciences

Relation between *in Utero* Arsenic Exposure and Birth Outcomes in a Cohort of Mothers and Their Newborns from New Hampshire

Diane Gilbert-Diamond,^{1,2*} Jennifer A. Emond,^{3*} Emily R. Baker,^{2,4} Susan A. Korrick,^{2,5,6} and Margaret R. Karagas^{1,2}

¹Department of Epidemiology, Geisel School of Medicine at Dartmouth, Hanover, New Hampshire, USA; ²Children's Environmental Health and Disease Prevention Center at Dartmouth, Hanover, New Hampshire, USA; ³Department of Biomedical Data Science, Geisel School of Medicine at Dartmouth, Hanover, New Hampshire, USA; ⁴Department of Obstetrics and Gynecology, Dartmouth-Hitchcock Medical Center, Lebanon, New Hampshire, USA; ⁵Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, New Hampshire, USA; ⁶Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, New Hampshire, USA. *Indicates equal contribution to this paper

Address correspondence to Diane Gilbert-Diamond, Hinman 7927, Geisel School of Medicine at Dartmouth, Hanover, New Hampshire 03755 USA. Telephone: (603) 653 3362. E-mail:

Diane.Gilbert-Diamond@Dartmouth.edu

Running title: Arsenic exposure and birth outcomes

Acknowledgments: We would like to thank the study participants and our sources of funding: National Institutes of Health (NIH) National Institute of Environmental Health Sciences P42ES007373, P01ES022832, and P42ES004940, NIH National Institute of General Medical

Sciences grants P20GM104416, Environmental Protection Agency RD83544201. No funding bodies had any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing financial interests: The authors declare they have no actual or potential competing financial interests.

Abstract

Background: Studies suggest that arsenic exposure influences birth outcomes, however findings are mixed.

Objective: We assessed *in utero* arsenic exposure in relation to birth outcomes and whether maternal pre-pregnancy weight and infant sex modified associations.

Methods: Among 706 mother-infant pairs exposed to low levels of arsenic through drinking water and diet, we assessed *in utero* arsenic exposure using maternal second trimester urinary arsenic, maternal pre-pregnancy weight through self-report, and birth outcomes from medical records.

Results: Median (IQ range) of total urinary arsenic (tAs; inorganic arsenic (iAs) + monomethylarsonic acid (MMA) + dimethylarsinic acid (DMA)) was 3.4 $\mu\text{g/L}$ (1.7 – 6.0). In adjusted linear models, each doubling of tAs was associated with a 0.10 cm decrease (95% CI: -0.19, -0.01) in head circumference. Results were similar for MMA and DMA. Ln(tAs) and Ln(DMA) were positively associated with birth length in infant males only; among males, each doubling of tAs was associated with a 0.28 cm increase (95% CI: 0.09, 0.46) birth length ($P_{\text{interaction}} = 0.04$). Results were similar for DMA. Additionally, arsenic exposure was inversely related to Ponderal Index and associations differed by maternal weight. Each Ln(tAs) doubling of tAs was associated with a 0.55 kg/m^3 lower (95% CI: -0.82, -0.28, $P < 0.001$) Ponderal Index for infants of overweight/obese, but not normal weight, mothers, ($P_{\text{interaction}} < 0.01$). Finally, there was a significant interaction between maternal weight status, infant sex, and arsenic exposure on birth weight ($P_{\text{interaction}} = 0.03$). In girls born of overweight/obese mothers, each doubling of tAs was associated with a 62.9 g decrease (95% CI: -111.6, -14.2) in birth weight, though the association was null in the other strata.

Conclusions: Low-level arsenic exposure may impact fetal growth, and the associations may be modified by maternal weight status and infant sex.

Introduction

Fetal development is a period marked by rapid cell division, differentiation and tissue growth and is therefore highly sensitive to the negative effects of toxic exposures. Arsenic is a naturally occurring chemical element found in bedrock and soil and it readily dissolves into surrounding groundwater that may be used for drinking (Smedley and Kinniburgh 2002). There is also growing awareness that people are exposed to arsenic through consumption of rice and other foods (Gilbert-Diamond et al. 2011; Meharg et al. 2009). Arsenic crosses the placenta (Concha et al. 1998), and experimental studies suggest that arsenic may have reproductive toxicity (Ahmed et al. 2011; Davey et al. 2007; Tsang et al. 2012), though the epidemiological evidence relating arsenic exposure to fetal growth has been mixed. Three prospective studies have found negative associations between arsenic exposure biomarkers and birth weight (Claus Henn et al. 2016; Laine et al. 2015; Rahman et al. 2009; Yang et al. 2003), and a third found a suggestive negative association (Ferreccio et al. 2003). In contrast, two large cross-sectional studies did not find any association between arsenic and birth weight (Kwok et al. 2006; Myers et al. 2010). Almost all of these studies were performed in arsenic endemic regions with generally high water arsenic (e.g., $>50 \mu\text{g/L}$) and few studies have examined arsenic-birth outcome relationships in areas with predominately lower levels of arsenic exposure (e.g., drinking water arsenic $<10 \mu\text{g/L}$).

After ingestion, iAs is methylated into monomethylarsonic acid (MMA^{V}) which is then reduced to monomethylarsonous acid (MMA^{III}) (Vahter 2002). In a second methylation step, MMA^{III} is converted to dimethylarsinic acid (DMA^{V}), which is then reduced to dimethylarsinous acid (DMA^{III}). This biotransformation of iAs is incomplete and iAs, MMA^{V} and DMA^{V} are all

excreted in the urine (Francesconi et al. 2002; Watanabe 2013). While the reduced methylated forms, MMA^{III} and DMA^{III}, have also been found in urine of people exposed to high levels of iAs (Mandal et al. 2001; Valenzuela et al. 2005), they have not been observed in urine samples from populations exposed to lower levels of arsenic (Meliker et al. 2012; Slotova and Gheorghiu 2006). MMA^V and DMA^V are generally considered less toxic than iAs (Gebel 2002; Moore et al. 1997); however, some animal studies suggest that MMA^{III} and DMA^{III} may be even more toxic than iAs (Styblo et al. 2000).

Weight status may impact the efficiency of arsenic metabolism (Gomez-Rubio et al. 2011; Yunus and Persson 2008), although findings have been mixed. For example, a study among 624 adult women in Southern US and Northern Mexico (Gomez-Rubio et al. 2011) where 68.9% of women were overweight or obese, documented that the efficiency of arsenic metabolism increased with maternal BMI. In contrast, a study among 442 Bangladeshi pregnant women (Yunus and Persson 2008) documented no statistical differences in arsenic methylation rates by maternal weight status during the first trimester; however, only 5.4% of the women in that sample were overweight (BMI >25 kg/m²). Given the importance of maternal weight status in fetal growth (Hull et al. 2008; King 2006; Sewell et al. 2006; Wu et al. 2004) and maternal weight status' potential role in arsenic metabolism, it is important to understand if maternal weight status modifies arsenic's association with fetal growth outcomes.

The aims of this study were to assess the associations between *in utero* arsenic exposure and birth size among a cohort of US pregnant women with low-level arsenic exposure from diet and drinking water, and to determine if maternal pre-pregnancy weight modified that association.

Methods

Source Population

The New Hampshire Birth Cohort Study (NHBCS) is a prospective study designed to examine the associations of arsenic exposure on fetal growth and development during early childhood (Gilbert-Diamond et al. 2011). In the state of New Hampshire, residents commonly use unregulated private water supplies, and it is estimated that more than one in ten homes have drinking water wells with water arsenic concentrations that exceed the Environmental Protection Agency's recommended maximum of 10 µg/L (Karagas et al. 1998). Pregnant women aged 18-45 years old between 24-28 weeks of gestation were recruited from prenatal clinics in New Hampshire beginning in January 2009. Eligibility criteria included English literacy, the use of a private, unregulated water system (e.g., private well) at home, and a singleton pregnancy. The New Hampshire Birth Cohort Study enrolled 1,033 pregnant women by September 30, 2013. At the time of the analysis there were 1,020 live singleton births. Of those, pre-pregnancy weight status was not available for 111 women, maternal 2nd trimester spot urine samples was not available for an additional 151 women, and newborn anthropometric data was not available for 27 newborns. Of the remaining 731 mother-child dyads, 25 were excluded for the mother having a pre-pregnancy BMI <18.5 kg/m². Thus the final sample size of the analysis was 706 dyads. When comparing women who were included in the analysis with the entire population-based cohort, there were no significant differences related to baseline characteristics, arsenic exposures or birth outcomes (data not shown). Participants provided informed consent and all study procedures were approved by the Internal Review Board at Dartmouth College.

Primary Outcomes: Measures of infant size at birth

Infant length, weight, head circumference, gestational age (weeks) and sex at birth were collected from a medical record review by trained study staff. Ponderal index at birth was computed as kg/m^3 and age- and sex-adjusted birth lengths and birth weight percentiles were computed per the methods of Fenton (Fenton et al. 2013). Age- and sex-adjusted birth weight percentiles were used to classify infants' size-for-gestational age, which included small ($\leq 10^{\text{th}}$ percentile) (SGA), average (11-89th percentile) (AGA) and large-for-gestational age ($\geq 90^{\text{th}}$ percentile) (LGA) (Lesseur et al. 2013; Wilhelm-Benartzi et al. 2012).

Primary Exposure: In Utero Arsenic Exposure

Using previously described methods (Gilbert-Diamond et al. 2011), we collected and analyzed maternal 2nd trimester urinary arsenic. Women provided a spot urine sample collected at approximately 24-28 weeks of gestation that were analyzed for iAs (As^{III} and As^{V}), MMA^{V} and DMA^{V} using a high-performance liquid chromatography (HPLC) ICP-MS system (Larsen et al. 1993). The limits of detection (LODs) ranged from 0.01 to 0.39 $\mu\text{g/L}$ for the individual arsenic species across the analysis batches. The average LOD for iAs was 0.15 $\mu\text{g/L}$, and the average LODs for MMA and DMA were both 0.06 $\mu\text{g/L}$. Measurements below the detection limit were set to half the LOD. Total urinary arsenic concentrations (tAs) was determined by summing those four metabolites and excluding arsenobetaine, an organoarsenic compound that is unmetabolized and is thought to be nontoxic (Francesconi et al. 2002). Inorganic arsenic, MMA^{V} , DMA^{V} and tAs concentrations were right-skewed and thus values were natural log-transformed for analysis. In addition, we divided MMA^{V} by iAs and DMA^{V} by MMA^{V} to calculate the primary and secondary methylation indices (PMI and SMI), respectively (Chen et

al. 2003; Gilbert-Diamond et al. 2013; Tseng 2009). The PMI and SMI were analyzed as continuous predictors. Urinary creatinine, a marker of urinary dilution, was measured using a calorimetric assay (Assay #500701; Cayman Chemical, Ann Arbor, MI).

In addition to providing urine samples, women also provided drinking water samples that were analyzed for arsenic as previously described (Gilbert-Diamond et al. 2011). As rice (Cascio et al. 2011; Gilbert-Diamond et al. 2011; Meharg et al. 2009) and seafood (Adams et al. 1994; Francesconi 2010) are also known sources of arsenic exposure in the U.S., women also reported their rice and seafood intake for the three days prior to the urine sample collection on a brief questionnaire.

Maternal Weight Status

Women were asked to report their usual weight when not pregnant as part of the enrollment questionnaire. Those values were combined with height abstracted from the medical records to compute pre-pregnancy BMI (kg/m^2). Pre-pregnancy weight status was categorized as normal weight ($18.5\text{--}25 \text{ kg/m}^2$), or overweight or obese ($\geq 25 \text{ kg/m}^2$). Twenty-five women who were classified as underweight ($<18.5 \text{ kg/m}^2$) before the current pregnancy were excluded from this analysis.

Other Measures

Demographics and lifestyle characteristics and medical history were collected via prenatal and post-partum questionnaires. Trained study staff also conducted a medical record review to collect results of gestational blood glucose testing. As part of routine prenatal care, women typically completed a non-fasting glucose challenge test (GCT) between 24–28 weeks

gestation, where blood was sampled 1 hour after ingestion of 50 g of glucose. For those with a 1-hour GCT result of 140-199 mg/dL an additional, fasting oral glucose tolerance test (OGTT) was performed, where blood was sampled hourly for 3 hours after ingesting 100 g of glucose.

Because women with a 1-hour GCT result ≥ 200 mg/dL were considered to have had gestational diabetes, they did not undergo a subsequent OGTT. For this analysis, results from both the GCT and OGTT were used to define maternal gestational hyperglycemia. Specifically, gestational diabetes mellitus (GDM) was defined as a 1-hour GCT result ≥ 200 mg/dL or failing the 1-hour GCT (140-199 mg/dL) with 2 or more high values on the OGTT based on the American Diabetes Association (ADA) criteria for a normal OGTT (American College of Obstetricians and Gynecologists 2013) of blood glucose ≤ 95 mg/dL at baseline, ≤ 180 mg/dL at 1 hour, ≤ 155 mg/dL at 2 hours, and ≤ 140 mg/dL at 3 hours. Impaired glucose tolerance (IGT) was defined as failing the 1-hour GCT (140-199 mg/dL) and having one high value on the OGTT. For two participants, results from 1-hour GCT were not available yet a diagnosis of GDM was included in the medical record. Therefore, data on IGT status was available for 680 participants, and data on GDM status was available for 682 participants.

Statistical Analyses

All analyses were completed using R: A Language and Environment for Statistical Computing, version 3.0.2 (R Foundation for Statistical Computing, Vienna, Austria). A threshold of $P < 0.05$ was used to define associations as statistically significant. Baseline characteristics, maternal urinary arsenic concentrations, infant gestational age, and infant birth size measures were summarized overall and compared by maternal pre-pregnancy weight status using Chi-

Square test for categorical measures, t-tests for normally distributed continuous measures and Wilcoxon-Rank-Sum tests for non-normally distributed continuous measures.

We assessed the linearity of each dose-response relationship between each arsenic exposure (ln-transformed) and each outcome using generalized additive models (GAMs). GAMs were fit using the R language gam function included in the mgcv package. In our analyses, the shape of dose-response relationship was estimated using thin plate splines (Wood 2004). The quasi-Akaike Information Criteria (qAIC) is an indicator of model fit, with lower values indicating a better-fitting model (Wood 2004). When considering the inclusion of a non-linear term in each model, the criteria of a change in qAIC of 10 or greater (Burnham & Anderson, 1998) was considered evidence of an improved model fit over a linear model. Per that criterion, we did not find evidence of any non-linear dose-response relationships between total arsenic (ln-transformed) and each outcome.

Models were fit in the overall sample and stratified by maternal pre-pregnancy weight status and infant sex. In all models, variables associated with maternal pre-pregnancy weight status or birth size at the $p < 0.10$ level in bivariate analyses were included as covariates. In this study, we chose to explore arsenic's association with birth anthropometric outcomes while controlling for gestational age and did not explore potential mediation of any associations by gestational age. The models for the head circumference, birth length, birth weight and Ponderal index outcomes were adjusted for maternal age at enrollment (continuous), maternal education (ordinal, with the five categories listed in Table 1 assigned values of 1–5, respectively), maternal parity (continuous), infant sex, and gestational age (continuous). Models for the gestational age outcome were similar except they did not adjust for gestational age. Models for the sex- and age-

specific birth length or weight percentiles were also similar but they did not adjust for gestational age or infant sex. Models with PMI and SMI as the main predictors were additionally adjusted for tAs. For each outcome, models for the overall sample also included maternal pre-pregnancy weight status (dichotomous: BMI < 25 kg/m², BMI ≥ 25 kg/m²) as a covariate.

Models with multiplicative interaction terms between the urinary As exposure and maternal pre-pregnancy weight status were used to formally assess modification by maternal pre-pregnancy weight status; a Wald test was used to determine statistical significance of the multiplicative terms. In addition, because other studies report potential sex-specific effects of arsenic exposure (Davis et al. 2015; Kippler et al. 2012; Pilsner et al. 2012), we conducted similar interaction analyses to test for effect modification by infant sex on the relationships between arsenic and birth outcomes. For birth outcomes that showed evidence of effect modification (either statistically significant or suggestive) by either maternal weight status or infant sex, we estimated the associations with each arsenic exposure stratified by both maternal weight status and infant sex. To test for the significance of a 3-way interaction between maternal weight status, infant sex and each arsenic exposure, we fit a multivariable regression with main effect terms, 2-way multiplicative interaction terms, and a 3-way multiplicative interaction term, adjusted for covariates; we assessed the statistical significance of each 3-way multiplicative interaction term using a Wald's test.

Several sensitivity analyses were completed to assess the robustness of the results. All multivariate regression models were repeated: 1) excluding women who reported any seafood intake within the two days prior to the urine sample collection, 2) excluding any women who reported any rice intake within the two days prior to the urine sample collection and 3) adjusted

for creatinine for the subset of women with urinary creatinine concentrations available. We also repeated the analysis with the head circumference outcome additionally adjusting for 1) vaginal or caesarean birth and 2) birth weight. Finally, because pre-pregnancy weight status is positively associated with risk of developing GDM (Torloni et al. 2009), we completed exploratory analyses to determine if there was a multiplicative interaction between maternal hyperglycemia and arsenic exposure on infant birth size using a Wald test to determine statistical significance of the interaction terms.

Results

Among the 706 included participants, 300 (42.5%) were overweight or obese before their current pregnancy and specifically 113 (16.0%) were obese. Table 1 presents baseline characteristics by maternal pre-pregnancy weight status. Three women were missing education data, 2 women were missing parity data, and 67 women were missing gestational weight gain data. In addition, 25 women were missing data on gestational diabetes mellitus (GDM) status, and 27 women were missing data on impaired glucose tolerance (IGT) and their 1-hour blood glucose screening. Women who were overweight or obese were younger, more likely to have a lower level of education and less likely to be nulliparous. Home water arsenic concentrations and any intake of seafood or rice did not differ significantly by pre-pregnancy weight status. Rates of IGT and GDM did not differ significantly by pre-pregnancy weight status, nor did mean screening blood glucose. Urinary iAs was correlated with MMA and DMA (Spearman's $\rho = 0.71$, 0.59 , respectively; both $P < 0.001$) and tAs was most strongly correlated with DMA (Spearman's $\rho = 0.98$, $P < 0.001$). Unadjusted concentrations of maternal urinary arsenic did not differ by pre-pregnancy weight status, nor did the PMI and SMI.

Infant birth characteristics by maternal pre-pregnancy weight status are presented in Table 1. Maternal overweight or obese weight status was associated with an earlier gestational age at birth, though there was no difference in the proportion of births before 37 weeks by weight status (5.4% overall, $P=0.43$). Overweight or obese status was also associated with a larger head circumference at birth. When considering age and sex adjusted size at birth (data not shown), the average (SD) birth length percentile was 57 (29) among all infants and did not differ by maternal weight status ($P = 0.12$). The average (SD) birth weight percentile was significantly higher for infants whose mothers were overweight or obese compared to normal weight pre-pregnancy (57 (27) vs 51 (26); $P=0.001$). Additionally, mothers who were overweight or obese gave birth to children with a 0.5 kg/m^3 higher Ponderal index, on average, though this difference was only marginally statistically significant ($P = 0.07$).

Results from the adjusted generalized additive models supported linear dose-response relationships between maternal urinary arsenic concentrations and gestational age, head circumference, birth length, birth weight, Ponderal Index, birth length percentile and birth weight percentile (data not shown). Table 2 presents the adjusted parameter estimates from a series of linear regression models fitting those infant birth characteristics on natural log transformed maternal urinary arsenic concentrations, overall and stratified by maternal pre-pregnancy weight status. Maternal urinary MMA and tAs were negatively associated with infant head circumference at birth. After holding all other covariates constant, each doubling of tAs was associated with a 0.10 cm decrease (95% CI: -0.19, -0.01, $P = 0.03$) in head circumference. The association was not modified by maternal weight status and did not change after further adjustment for delivery mode and/or birth weight (see Supplemental Material, Table S1). Results were also consistent in unadjusted models (data not shown). The negative associations between

maternal urinary iAs and DMA and head circumference were similar in magnitude to those for MMA and tAs, however they did not reach statistical significance.

Maternal urinary DMA and tAs were positively associated with infant birth length. After holding all other covariates constant, each doubling of tAs was associated with a 0.15 cm increase (95% CI: 0.01, 0.28, $P = 0.04$) in birth length. While the estimates of association were larger and only statistically significant in the overweight/obese stratum, the statistical test for interaction by maternal weight status did not reach statistical significance. The results for birth length percentile were qualitatively similar to those for birth length (see Supplemental Material, Table S2); birth length percentile was positively associated with Ln(DMA) and Ln(tAs) in the stratum of overweight/obese women, though only the Ln(DMA) association was statistically significant.

Maternal urinary arsenic was inversely associated with infant adiposity measured as Ponderal Index and the associations appeared limited to women who were overweight or obese pre-pregnancy. Among children of those women, each doubling of tAs was associated with a 0.55 kg/m³ lower (95% CI: -0.82, -0.28, $P < 0.001$) Ponderal Index. Additionally among those women, decreases in mean Ponderal index also were observed with increases in urinary iAs, MMA and DMA. In contrast, there were not any statistically significant associations between maternal arsenic concentrations and infant Ponderal Index among normal weight women.

Maternal urinary arsenic was not significantly associated with gestational age, absolute birth weight (Table 2), or birth weight percentiles (Supplemental Material, Table S2). The PMI and SMI were not significantly associated with gestational age or any birth anthropometric measures (data not shown).

In analyses that examined effect modification by infant sex (Supplemental Material, Table S3), arsenic exposure was positively and significantly related to birth length in male infants though showed no association in females (P for interaction for $\text{Ln}(\text{tAs}) = 0.04$); among male infants, each doubling of tAs was associated with a 0.28 cm increase (95% CI: 0.09, 0.46, $P < 0.01$) birth length. Results were similar for DMA and MMA. There was also a suggestive interaction between arsenic exposure and infant sex on the birth weight outcome; arsenic appeared negatively associated with birth weight in girls and positively associated with birth weight in boys, though none of the point estimates were statistically significant, nor were any of the interaction tests.

In analyses stratified by both maternal weight status and infant sex (Table 3), the positive association between arsenic exposure and infant length appeared limited to the stratum of boys born of overweight/obese mothers, though the test for 3-way interaction was only marginally significant ($P = 0.13$). In the stratum of boys born of overweight/obese mothers, each doubling of tAs was associated with a 0.51 increase (95% CI: 0.21, 0.82) in birth length. When looking at the birth weight outcome, there was a strong, statistically significant negative association between arsenic exposure and birth weight in the stratum of girls born of overweight/obese moms, while associations in the other strata were largely positive and not statistically significant; the test for 3-way interaction was significant ($P = 0.03$). In the stratum of girls born of overweight/obese mothers, each doubling of tAs was associated with a 62.9 g decrease (95% CI: -111.6, -14.2) in birth weight.

Results were consistent in analyses limited to the subset of women who did not consume seafood ($N=594$) or rice ($N=543$) in the previous two days, with additional adjustment for

creatinine (data available for 520 participants) (data not shown), or with additional adjustment for gestational weight gain (data available for 639 participants) (data not shown). Also, we did not find evidence that the results of the maternal glucose challenge test or maternal hyperglycemia status modified the association between maternal urinary tAs and infant Ponderal Index (data not shown).

Discussion

In this U.S. sample of women and their newborn infants, we found that 2nd trimester maternal urinary arsenic concentration, a measure of fetal *in utero* arsenic exposure, was negatively associated with infant head circumference, positively associated with infant length and, negatively associated with adiposity at birth. We further observed that the associations between arsenic exposure and birth length were only observed in males, and the associations with adiposity were only observed in the overweight/obese stratum and not in the normal weight stratum. Finally, we observed a significant 3-way interaction between maternal weight status, infant sex, and arsenic exposure on the birth weight outcome; specifically, arsenic exposure is related to lower birth weight in females born to overweight/obese mothers, but the association appears null, or possibly in the opposite direction, for all males and for females born of healthy weight mothers.

Our findings relating arsenic exposure and head circumference at birth agree with those from a large prospective analysis by Rahman *et al.* (2009). In that study of 805 Bangladeshi mothers with urinary arsenic in the lower range of exposure (<100 µg/L) and their children, 1 µg/L of maternal tAs (average from samples taken around gestational weeks 8 and 30) was associated with a 0.05 mm (SE=0.03, $P = 0.04$) lower newborn head circumference after

adjustment for maternal BMI and socioeconomic status. An analysis of ultrasound measurements from the Bangladeshi cohort (n=1929) found that arsenic may influence head circumference before the 3rd trimester and in a sex-specific manner (Kippler et al. 2012); maternal urinary arsenic was negatively associated with ultrasound measures of head circumference in males and positively associated with head circumference in females. Our previous analysis of 223 ultrasound reports from our cohort also suggests that this arsenic-related reduced head circumference may be present as early as the second trimester, though we observed a stronger negative association in female fetuses (Davis et al. 2015). In the present study, we did not observe any significant effect modification of the associations between arsenic and head circumference by infant sex.

In contrast to our results, the Rahman study did not observe an association between arsenic exposure and birth length. The Rahman study did observe a negative association between maternal urinary arsenic and birth weight; each 1 µg/L of urinary tAs was associated with an adjusted -1.68 g (SE=0.62, $P < 0.01$) difference in birth weight when examining infants born of mothers with urinary arsenic concentrations <100 µg/L. In contrast, we only observed a negative association between arsenic and birth weight in girls born of overweight/obese mothers.

Three other prospective studies have examined maternal urinary arsenic in relation to birth outcomes (Chou et al. 2014; Laine et al. 2015; Shirai et al. 2010). A study of 299 maternal-child pairs from a non-arsenic-endemic region of Taiwan (Chou et al. 2014) and a study of 78 low-exposure maternal-child pairs from Tokyo (Shirai et al. 2010), did not find any associations between maternal urinary arsenic and head circumference, birth length or birth weight. In a study of 200 maternal-child pairs from an arsenic-endemic region in Mexico, maternal urinary MMA

was negatively associated with birth weight and gestational age, and iAs was negatively associated with birth length and gestational age (Laine et al. 2014). In a small study that compared 19 mothers from a city with high water arsenic to 19 mothers from a city with low water arsenic in Chile, there were no significant urinary arsenic-birth weight associations in (Ferreccio et al. 2003).

The results are also mixed from other studies that relied on maternal hair (Breton and Afroz 2007), maternal blood (Claus Henn et al. 2016; Guan et al. 2012), infant meconium (Vall et al. 2012) and drinking water (Ferreccio et al. 2003; Kwok et al. 2006; Myers et al. 2010; Yang et al. 2003) arsenic levels to examine the association between arsenic and birth weight (Breton and Afroz 2007; Ferreccio et al. 2003; Guan et al. 2012; Kwok et al. 2006; Myers et al. 2010; Vall et al. 2012; Yang et al. 2003, Claus Henn et al. 2016); length (Guan et al. 2012; Vall et al. 2012, Claus Henn et al. 2016); and head circumference (Guan et al. 2012; Vall et al. 2012, Claus Henn et al. 2016).

Several population characteristics could lead to the heterogeneity of study findings. Our research differs from the bulk of previous research on arsenic and fetal growth because of the relatively low levels of arsenic exposure in our population; for example, participants in the Rahman study had a median concentration of 95 $\mu\text{g/L}$ of total urinary arsenic compared to our median of 3.4 $\mu\text{g/L}$. In addition, almost half of our study population was overweight or obese. In contrast, only 5% of the mothers in the Rahman study were overweight or obese and 30% were underweight. We were unable to explore the relation between arsenic and fetal growth in underweight women, as only 25 women in our sample met this criterion. In addition to weight status, other nutritional factors such as folate (Gamble et al. 2007; Hall et al. 2009) and protein

intake (Heck et al. 2009; Milton et al. 2007) could modify the associations between arsenic and fetal growth; thus nutritional factors may lead to heterogeneity in study findings and should be explored in future research.

Fetal growth is a complicated process that is affected by several factors including maternal nutrition, genetics, and toxic exposures (Kramer 1987). While the mechanism by which arsenic might impact fetal growth is unknown, several possibilities have been proposed. Our previous work suggests that increased maternal arsenic exposure may lead to increased expression of the arsenic transporter AQP9 in placental tissue, which, in turn, may decrease expression of the adipose tissue-derived ENPP2 (Fei et al. 2013), a phospholipase that is purported to regulate adipose tissue growth (Nishimura et al. 2014). In addition, there is evidence that arsenic increases oxidative stress (Xu et al. 2008) and inflammatory processes (Ahmed et al. 2011; Fry et al. 2007), factors associated with growth restriction (Biri et al. 2007; Challis et al. 2009). Given that excess adiposity is also known to increase oxidative stress and systemic inflammation (Vincent and Taylor 2006), it is feasible that arsenic exposure may differentially impact fetal growth depending on maternal weight status. In addition, arsenic could influence fetal growth through other mechanisms such as endocrine disruption (Davey et al. 2007; Jack E Bodwell et al. 2004), epigenetic modification (Koestler et al. 2013), and/or altered transcription (Kumagai and Sumi 2007) and it is plausible that effects could differ by infant sex.

Opposite effects of *in utero* arsenic exposure on fetal length at lower and higher exposure doses is plausible given the findings of an *in vitro* study of arsenic trioxide (ATO) and the growth of osteoblasts—the specialized cells in the bone that secrete collagen and a calcium- and phosphate-containing mineral to form the bone matrix (Xu et al. 2014). In that study a low-dose

(0.25, 0.5 and 1 μ M) of ATO, an oxide of inorganic arsenic, greatly enhanced the viability of cultured osteoblasts and promoted collagen synthesis and secretion, while a high-dose (5, 10, and 20 μ M) of ATO led to a significant reduction in osteoblast viability and induced osteoblast apoptosis. While we did not observe an inverse association between arsenic and birth length at any part of the dose-response curves in the stratified or total samples, this may be due to the relatively low range of arsenic exposure in our population.

Alterations of fetal growth related to *in utero* arsenic exposure may have important public health implications. Head circumference closely correlates to brain volume (Cooke et al. 1977; Wickett et al. 2000) and some research suggests that head circumference at birth is associated with later intellectual function (Gale et al. 2006). Additionally, the persistence of the negative association between arsenic and head circumference after adjusting for birth weight suggests that the reduction in head circumference is independent of birth size as has been observed with other toxicant exposure (Fein et al. 1984; Hernandez-Avila et al. 2002). Thus, our findings of increased arsenic exposure related to decreased head circumference are consistent with laboratory studies that find that arsenic has neurotoxic effects (Chaudhuri et al. 1999), and epidemiological studies that suggest that arsenic exposure is related to decreased intellectual function (Hamadani et al. 2011; Roy et al. 2011). Other studies suggest that postnatal head growth is also important in determining later cognitive function (Gale et al. 2004; Smithers et al. 2013) and so it is important to further explore arsenic's relationship with postnatal head circumference growth and ultimately cognitive function in future studies.

The influence of fetal growth on later risk of adiposity (Curhan et al. 1996a; 1996b) and metabolic disease (Barker et al. 1989b; 1989a) is well recognized. Studies suggest a U-shaped

relationship with both high birth weight/adiposity (Eriksson et al. 2001) and low birth weight and adiposity (Barker et al. 1993; Bhargava et al. 2004) related to later adiposity and metabolic disease. Studies additionally suggest that low size at birth may be related to accelerated weight gain (catch up growth) in infancy (Barker et al. 1993; Bhargava et al. 2004) which is in turn related to later obesity (Hack et al. 1984; Ong et al. 2000; Victora et al. 2001). Interestingly, our observed negative association between arsenic exposure and Ponderal Index in the overweight/obese stratum seems to be driven by longer birth length in boys, but by lower birth weight in girls. Such possible mechanistic differences in adiposity at birth could have important ramifications for later growth trajectories. Ongoing follow up of our longitudinal cohort will allow us to examine weight and adiposity trajectories in early infancy and, ultimately childhood overweight or obesity status, in relation to *in utero* arsenic exposure in our population. We will also be able to explore whether early childhood arsenic exposure is related to early childhood growth as has been done in more highly exposed populations (Gardner et al. 2013). A study of 1,505 mother-infant pairs in rural Bangladesh found an inverse relationship between children's weight, height, and growth velocity and concurrent arsenic exposure at 5 years of age (Gardner et al. 2013).

Unlike the majority of epidemiological studies on arsenic exposure, this population-based prospective study explores arsenic exposures at levels commonly encountered in the U.S. through water and food sources. Our study is strengthened by using sensitive biomarkers of *in utero* exposure to both inorganic arsenic and its metabolites as well as clinically measured anthropometry at birth. While we only have a single measure of maternal urinary arsenic during pregnancy, a study in American Indian communities suggest that urinary arsenic is relatively stable over time (Navas-Acien et al. 2009). Our study was also limited by our use of self-

reported usual pre-pregnancy weight, which is likely slightly under-reported (Holland et al. 2013; Russell et al. 2013), which may have led to some women who were overweight/obese being mis-classified as healthy weight. In this analysis, we only explored arsenic species and did not examine arsenic's correlation with other metals so we cannot rule out the potential for confounding by those factors. In addition, we do not adjust for micronutrient status, which may relate to arsenic metabolism; however, we believe it is unlikely that this would have substantially influenced our results as all women received prenatal care at study clinics and approximately 92% of women in the cohort report taking prenatal vitamins. A further limitation is that we were underpowered to explore arsenic's association specifically with risk for being born SGA or LGA, given the small number of infants born with these outcomes.

Our study, in a U.S. cohort of mothers and infants, suggest that arsenic exposure, even at low levels, may impact fetal growth, and that associations may be modified by maternal weight status and infant sex. The health implications of our findings will be elucidated by future longitudinal studies in our cohort and others that explore the impact of prenatal and postnatal arsenic exposure on growth and neurodevelopmental outcomes.

References

- Adams MA, Gunderson EL, Bolger PM. 1994. Dietary intake and hazards of arsenic. In *Arsenic Exposure and Health* (W.R. Chapell, C.O. Abernathy, and C.R. Cotherneds.), Northwood, U.K.
- Ahmed S, Mahabbat-e Khoda S, Rekha RS, Ameer SS, Moore S, Raqib R. 2011. Arsenic-associated oxidative stress, inflammation, and immune disruption in human placenta and cord blood. *Environ. Health Perspect.* 119:258–264; doi:10.1289/ehp.1002086.
- American College of Obstetricians and Gynecologists. 2013. ACOG Practice Bulletin no. 137: Clinical Management Guidelines for Obstetrician-Gynecologists. Gestational diabetes. *Obstet Gynecol* 122: 406–416.
- Barker DJ, Hales CN, Osmond C, Phipps K, Clark PM. 1993. Type 2 (non-insulin-dependent) diabetes mellitus, hypertension and hyperlipidaemia (syndrome X): relation to reduced fetal growth. *Diabetologia* 36: 62–67.
- Barker DJ, Osmond C, Golding J, Kuh D, Wadsworth ME. 1989a. Growth in utero, blood pressure in childhood and adult life, and mortality from cardiovascular disease. *BMJ* 298:564–567; doi:10.1136/bmj.298.6673.564.
- Barker DJP, Osmond C, Winter PD, Margetts B. 1989b. Weight in infancy and death from ischaemic Heart Disease. *The Lancet* 334:577–580; doi:10.1016/S0140-6736(89)90710-1.
- Bhargava SK, Sachdev HS, Fall CHD, Osmond C, Lakshmy R, Barker DJP, et al. 2004. Relation of serial changes in childhood body-mass index to impaired glucose tolerance in young adulthood. *N. Engl. J. Med.* 350:865–875; doi:10.1056/NEJMoa035698.
- Biri A, Bozkurt N, Turp A, Kavutcu M, Himmetoglu O, Durak I. 2007. Role of oxidative stress in intrauterine growth restriction. *Gynecol. Obstet. Invest.* 64:187–192; doi:10.1159/000106488.
- Breton CV, Afroz S. 2007. Maternal arsenic exposure associated with low birth weight in Bangladesh. *J. Occup. Environ. Med.* 49:1097–104; doi:10.1097/JOM.0b013e3181566ba0.
- Cascio C, Raab A, Jenkins RO, Feldmann J, Meharg AA, Haris PI. 2011. The impact of a rice based diet on urinary arsenic. *J. Environ. Monit.* 13:257–265; doi:10.1039/c0em00482k.
- Challis JR, Lockwood CJ, Myatt L, Norman JE, Strauss JF III, Petraglia F. 2009. Inflammation and Pregnancy. *Reproductive Sciences* 16:206–215; doi:10.1177/1933719108329095.

- Chaudhuri AN, Basu S, Chattopadhyay S. 1999. Effect of high arsenic content in drinking water on rat brain. *Indian J Biochem and Biophys* 36: 51–54.
- Chen Y-C, Su H-JJ, Guo Y-LL, Hsueh Y-M, Smith TJ, Ryan LM, et al. 2003. Arsenic methylation and bladder cancer risk in Taiwan. *Cancer Causes Control* 14: 303–310.
- Chou W-C, Chung Y-T, Chen H-Y, Wang C-J, Ying T-H, Chuang C-Y, et al. 2014. Maternal arsenic exposure and DNA damage biomarkers, and the associations with birth outcomes in a general population from Taiwan. *J. Melikered. PLoS ONE* 9:e86398; doi:10.1371/journal.pone.0086398.
- Claus Henn B, Ettinger AS, Hopkins MR, Jim R, Amarasiriwardena C, Christiani DC, et al. 2016. Prenatal Arsenic Exposure and Birth Outcomes among a Population Residing near a Mining-Related Superfund Site. *Environ. Health Perspect.*; doi:10.1289/ehp.1510070.
- Concha G, Vogler G, Lezcano D, Nermell B, Vahter M. 1998. Exposure to inorganic arsenic metabolites during early human development. *Toxicol. Sci.* 44:185–190; doi:10.1006/toxs.1998.2486.
- Cooke RWI, Lucas A, Yudkin PL, Pryse-Davies J. 1977. Head circumference as an index of brain weight in the fetus and newborn. *Early Human Development* 1:145–149; doi:10.1016/0378-3782(77)90015-9.
- Curhan GC, Chertow GM, Willett WC, Spiegelman D, Colditz GA, Manson JE, et al. 1996a. Birth weight and adult hypertension and obesity in women. *Circulation* 94:1310–1315; doi:10.1161/01.CIR.94.6.1310.
- Curhan GC, Willett WC, Rimm EB, Spiegelman D, Ascherio AL, Stampfer MJ. 1996b. Birth weight and adult hypertension, diabetes mellitus, and obesity in US men. *Circulation* 94:3246–3250; doi:10.1161/01.CIR.94.12.3246.
- Davey JC, Bodwell JE, Gosse JA, Hamilton JW. 2007. Arsenic as an Endocrine Disruptor: Effects of Arsenic on Estrogen Receptor-Mediated Gene Expression In Vivo and in Cell Culture. *Toxicol. Sci.* 98:75–86; doi:10.1093/toxsci/kfm013.
- Davis MA, Higgins J, Li Z, Gilbert-Diamond D, Baker ER, Das A, et al. 2015. Preliminary analysis of in utero low-level arsenic exposure and fetal growth using biometric measurements extracted from fetal ultrasound reports. *Environmental Health* 14:12; doi:10.1186/1476-069X-14-12.
- Eriksson J, Forsén T, Tuomilehto J, Osmond C, Barker D. 2001. Size at birth, childhood growth and obesity in adult life. *Int. J. Obes. Relat. Metab. Disord.* 25:735–740;

doi:10.1038/sj.ijo.0801602.

- Fei DL, Koestler DC, Li Z, Giambelli C, Sanchez-Mejias A, Gosse JA, et al. 2013. Association between In Utero arsenic exposure, placental gene expression, and infant birth weight: a US birth cohort study. *Environ Health* 12:58; doi:10.1186/1476-069X-12-58.
- Fein GG, Jacobson JL, Jacobson SW. 1984. Prenatal exposure to polychlorinated biphenyls: effects on birth size and gestational age. *J Pediatr* 105: 315–320.
- Fenton TR, Nasser R, Eliasziw M, Kim JH, Bilan D, Sauve R. 2013. Validating the weight gain of preterm infants between the reference growth curve of the fetus and the term infant. *BMC Pediatr* 13:92; doi:10.1186/1471-2431-13-92.
- Ferreccio C, Browning SR, Huang B, Peralta C, Gibb H, Hertz-Picciotto I. 2003. Arsenic exposure from drinking water and birth weight. *Epidemiology* 14:593–602; doi:10.1097/01.ede.0000072104.65240.69.
- Francesconi KA. 2010. Arsenic species in seafood: Origin and human health implications. *Pure and Applied Chemistry* 82:373–381; doi:10.1351/PAC-CON-09-07-01.
- Francesconi KA, Tanggaar R, McKenzie CJ. 2002. Arsenic metabolites in human urine after ingestion of an arsenosugar. *Clin. Chem.* 48: 92–101.
- Fry RC, Navasumrit P, Valiathan C, Svensson JP, Hogan BJ, Luo M, et al. 2007. Activation of Inflammation/NF- κ B Signaling in Infants Born to Arsenic-Exposed Mothers. *PLoS Genet.* 3:e207; doi:10.1371/journal.pgen.0030207.
- Gale CR, O'Callaghan FJ, Bredow M, Martyn CN, Avon Longitudinal Study of Parents and Children Study Team. 2006. The influence of head growth in fetal life, infancy, and childhood on intelligence at the ages of 4 and 8 years. *PEDIATRICS* 118:1486–1492; doi:10.1542/peds.2005-2629.
- Gale CR, O'Callaghan FJ, Godfrey KM, Law CM, Martyn CN. 2004. Critical periods of brain growth and cognitive function in children. *Brain* 127:321–329; doi:10.1093/brain/awh034.
- Gamble MV, Liu X, Slavkovich V, Pisner JR, Ilievski V, Factor-Litvak P, et al. 2007. Folic acid supplementation lowers blood arsenic. *Am J of Clin Nutr* 86: 1202–1209.
- Gardner RM, Kippler M, Tofail F, Bottai M, Hamadani J, Grandér M, et al. 2013. Environmental exposure to metals and children's growth to age 5 years: a prospective cohort study. *Am. J. Epidemiol.* 177:1356–1367; doi:10.1093/aje/kws437.

- Gebel TW. 2002. Arsenic methylation is a process of detoxification through accelerated excretion. *Int J Hyg Environ Health* 205:505–508; doi:10.1078/1438-4639-00177.
- Gilbert-Diamond D, Cottingham KL, Gruber JF, Punshon T, Sayarath V, Gandolfi AJ, et al. 2011. Rice consumption contributes to arsenic exposure in US women. *Proc. Natl. Acad. Sci. U.S.A.* 108:20656–20660; doi:10.1073/pnas.1109127108.
- Gilbert-Diamond D, Li Z, Perry AE, Spencer SK, Gandolfi AJ, Karagas MR. 2013. A population-based case-control study of urinary arsenic species and squamous cell carcinoma in New Hampshire, USA. *Environ. Health Perspect.* 121:1154–1160; doi:10.1289/ehp.1206178.
- Gomez-Rubio P, Roberge J, Arendell L, Harris RB, O'Rourke MK, Chen Z, et al. 2011. Association between body mass index and arsenic methylation efficiency in adult women from southwest U.S. and northwest Mexico. *Toxicol. Appl. Pharmacol.* 252:176–182; doi:10.1016/j.taap.2011.02.007.
- Guan H, Piao F, Zhang X, Li X, Li Q. 2012. Prenatal Exposure to Arsenic and Its Effects on Fetal Development in the General Population of Dalian. *Biol Trace Elem Res* 149:10–15; doi:10.1007/s12011-012-9396-7.
- Hack M, Merikatz IR, McGrath SK, Jones PK, Fanaroff AA. 1984. Catch-up growth in very-low-birth-weight infants. Clinical correlates. *Am. J. Dis. Child.* 138: 370–375.
- Hall MN, Liu X, Slavkovich V, Ilievski V, Pilsner JR, Alam S, et al. 2009. Folate, Cobalamin, Cysteine, Homocysteine, and Arsenic Metabolism among Children in Bangladesh. *Environ. Health Perspect.* 117:825–831; doi:10.1289/ehp.0800164.
- Hamadani JD, Tofail F, Nermell B, Gardner R, Shiraji S, Bottai M, et al. 2011. Critical windows of exposure for arsenic-associated impairment of cognitive function in pre-school girls and boys: a population-based cohort study. *Int J Epidemiol* 40:1593–1604; doi:10.1093/ije/dyr176.
- Heck JE, Nieves JW, Chen Y, Parvez F, Brandt-Rauf PW, Graziano JH, et al. 2009. Dietary intake of methionine, cysteine, and protein and urinary arsenic excretion in Bangladesh. *Environ. Health Perspect.* 117:99–104; doi:10.1289/ehp.11589.
- Hernandez-Avila M, Hernandez AV, Peterson KE, Gonzalez-Cossio T, Sanin LH, Aro A, et al. 2002. Effect of maternal bone lead on length and head circumference of newborns and 1-month-old infants. *Arch Environ Health* 57:482–488; doi:10.1080/00039890209601441.
- Holland E, Moore Simas TA, Doyle Curiale DK, Liao X, Waring ME. 2013. Self-reported pre-

- pregnancy weight versus weight measured at first prenatal visit: effects on categorization of pre-pregnancy body mass index. *Matern Child Health J* 17:1872–1878; doi:10.1007/s10995-012-1210-9.
- Hull HR, Dinger MK, Knehans AW, Thompson DM, Fields DA. 2008. Impact of maternal body mass index on neonate birthweight and body composition. *Am. J. Obstet. Gynecol.* 198:416.e1–6; doi:10.1016/j.ajog.2007.10.796.
- Jack E Bodwell, Lauren A Kingsley A, Hamilton JW. 2004. Arsenic at Very Low Concentrations Alters Glucocorticoid Receptor (GR)-Mediated Gene Activation but Not GR-Mediated Gene Repression: Complex Dose–Response Effects Are Closely Correlated with Levels of Activated GR and Require a Functional GR DNA Binding Domain. *Chem. Res. Toxicol.* 17:1064–1076; doi:10.1021/tx0499113.
- Karagas MR, Tosteson TD, Blum J, Morris JS, Baron JA, Klaue B. 1998. Design of an epidemiologic study of drinking water arsenic exposure and skin and bladder cancer risk in a U.S. population. *Environ. Health Perspect.* 106 Suppl 4: 1047–1050.
- Kile ML, Hoffman E, Hsueh Y-M, Afroz S, Quamruzzaman Q, Rahman M, et al. 2009. Variability in biomarkers of arsenic exposure and metabolism in adults over time. *Environ. Health Perspect.* 117:455–460; doi:10.1289/ehp.11251.
- King JC. 2006. Maternal obesity, metabolism, and pregnancy outcomes. *Annu. Rev. Nutr.* 26:271–291; doi:10.1146/annurev.nutr.24.012003.132249.
- Kippler M, Wagatsuma Y, Rahman A, Nermell B, Persson LA, Raqib R, et al. 2012. Environmental exposure to arsenic and cadmium during pregnancy and fetal size: A longitudinal study in rural Bangladesh. *Reproductive Toxicology* 34:504–511; doi:10.1016/j.reprotox.2012.08.002.
- Koestler DC, Avissar-Whiting M, Houseman EA, Karagas MR, Marsit CJ. 2013. Differential DNA methylation in umbilical cord blood of infants exposed to low levels of arsenic in utero. *Environ. Health Perspect.* 121:971–977; doi:10.1289/ehp.1205925.
- Kramer MS. 1987. Determinants of low birth weight: methodological assessment and meta-analysis. *Bulletin of the World Health Organization* 65: 663–737.
- Kumagai Y, Sumi D. 2007. Arsenic: signal transduction, transcription factor, and biotransformation involved in cellular response and toxicity. *Annu. Rev. Pharmacol. Toxicol.* 47:243–262; doi:10.1146/annurev.pharmtox.47.120505.105144.
- Kwok RK, Kaufmann RB, Jakariya M. 2006. Arsenic in drinking-water and reproductive health

outcomes: a study of participants in the Bangladesh Integrated Nutrition Programme. *J Health Popul Nutr* 24: 190–205.

Laine JE, Bailey KA, Rubio-Andrade M, Olshan AF, Smeester L, Drobná Z, et al. 2015.

Maternal arsenic exposure, arsenic methylation efficiency, and birth outcomes in the Biomarkers of Exposure to ARsenic (BEAR) pregnancy cohort in Mexico. *Environ. Health Perspect.* 123:186–192; doi:10.1289/ehp.1307476.

Laine JE, Bailey KA, Rubio-Andrade M, Olshan AF, Smeester L, Drobná Z, et al. 2014.

Maternal Arsenic Exposure, Arsenic Methylation Efficiency, and Birth Outcomes in the Biomarkers of Exposure to ARsenic (BEAR) Pregnancy Cohort in Mexico. *Environ. Health Perspect.* 1–7; doi:10.1289/ehp.1307476.

Larsen EH, Pritzl G, Hansen SH. 1993. Speciation of 8 Arsenic Compounds in Human Urine by High-Performance Liquid-Chromatography with Inductively-Coupled Plasma-Mass Spectrometric Detection Using Antimonate for Internal Chromatographic Standardization. *J. Anal. At. Spectrom.* 8:557–563; doi:10.1039/ja9930800557.

Lesseur C, Armstrong DA, Paquette AG, Koestler DC, Padbury JF, Marsit CJ. 2013. Tissue-specific Leptin promoter DNA methylation is associated with maternal and infant perinatal factors. *Molecular and Cellular Endocrinology* 381:160–167; doi:10.1016/j.mce.2013.07.024.

Mandal BK, Ogra Y, Suzuki KT. 2001. Identification of dimethylarsinous and monomethylarsonous acids in human urine of the arsenic-affected areas in West Bengal, India. *Chem. Res. Toxicol.* 14:371–378; doi:10.1021/tx000246h.

Meharg AA, Williams PN, Adomako E. 2009. Geographical variation in total and inorganic arsenic content of polished (white) rice. *Environ. Sci. Technol.* 43: 1612–1617.

Meliker JR, Meeker JD, Slotnick MJ. 2012. Urinary arsenic species, toenail arsenic, and arsenic intake estimates in a Michigan population with low levels of arsenic in drinking water. *J Expo Sci Environ Epidemiol* 22:182–190; doi:10.1038/jes.2011.27.

Milton AH, Hasan Z, Shahidullah SM, Sharmin S, Jakariya MD, Dear K, et al. 2007. Association between nutritional status and arsenicosis due to chronic arsenic exposure in Bangladesh. *International Journal of Environmental Health Research* 14:99–108; doi:10.1080/0960312042000209516.

Moore MM, Harrington-Brock K, Doerr CL. 1997. Relative genotoxic potency of arsenic and its methylated metabolites. *Mutation Research/Reviews in Mutation Research* 386:279–290; doi:10.1016/S1383-5742(97)00003-3.

- Myers SL, Lobdell DT, Liu Z, Xia Y, Ren H, Li Y, et al. 2010. Maternal drinking water arsenic exposure and perinatal outcomes in inner Mongolia, China. *J Epidemiol Community Health* 64:325–329; doi:10.1136/jech.2008.084392.
- Navas-Acien A, Umans JG, Howard BV, Francesconi KA, Crainiceanu CM, Silbergeld EK, et al. 2009. Urine arsenic concentrations and species excretion patterns in American Indian communities over a 10-year period: the Strong Heart Study. *Environ. Health Perspect.* 117:1428–1433; doi:10.1289/ehp.0800509.
- Nishimura S, Nagasaki M, Okudaira S, Aoki J, Ohmori T, Ohkawa R, et al. 2014. ENPP2 contributes to adipose tissue expansion and insulin resistance in diet-induced obesity. *Diabetes* 63:4154–4164; doi:10.2337/db13-1694.
- Ong KK, Ahmed ML, Emmett PM, Preece MA, Dunger DB. 2000. Association between postnatal catch-up growth and obesity in childhood: prospective cohort study. *BMJ* 320: 967–971.
- Pilsner JR, Hall MN, Liu X, Ilievski V, Slavkovich V, Levy D, et al. 2012. Influence of prenatal arsenic exposure and newborn sex on global methylation of cord blood DNA. *C. Oudejansed. PLoS ONE* 7:e37147; doi:10.1371/journal.pone.0037147.
- Rahman A, Vahter M, Smith AH, Nermell B, Yunus M, Arifeen SE, et al. 2009. Arsenic exposure during pregnancy and size at birth: a prospective cohort study in Bangladesh. *Am. J. Epidemiol.* 169:304–312; doi:10.1093/aje/kwn332.
- Roy A, Kordas K, Lopez P, Rosado JL, Cebrian ME, Vargas GG, et al. 2011. Association between arsenic exposure and behavior among first-graders from Torreón, Mexico. *Environ. Res.* 111:670–676; doi:10.1016/j.envres.2011.03.003.
- Russell A, Gillespie S, Satya S, Gaudet LM. 2013. Assessing the accuracy of pregnant women in recalling pre-pregnancy weight and gestational weight gain. *J Obstet Gynaecol Can* 35: 802–809.
- Sewell MF, Huston-Presley L, Super DM, Catalano P. 2006. Increased neonatal fat mass, not lean body mass, is associated with maternal obesity. *Am. J. Obstet. Gynecol.* 195:1100–1103; doi:10.1016/j.ajog.2006.06.014.
- Shirai S, Suzuki Y, Yoshinaga J, Mizumoto Y. 2010. Maternal exposure to low-level heavy metals during pregnancy and birth size. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 45:1468–1474; doi:10.1080/10934529.2010.500942.
- Slotova K, Gheorghiu E. 2006. Arsenic exposure in Hungary, Romania and Slovakia. *J. Environ.*

- Monit. 8:203–208; doi:10.1039/b513206a.
- Smedley PL, Kinniburgh DG. 2002. A review of the source, behaviour and distribution of arsenic in natural waters. *Applied Geochemistry* 17:517–568; doi:10.1016/S0883-2927(02)00018-5.
- Smithers LG, Lynch JW, Yang S, Dahhou M, Kramer MS. 2013. Impact of neonatal growth on IQ and behavior at early school age. *PEDIATRICS* 132:e53–60; doi:10.1542/peds.2012-3497.
- Styblo M, Del Razo LM, Vega L, Germolec DR, LeCluyse EL, Hamilton GA, et al. 2000. Comparative toxicity of trivalent and pentavalent inorganic and methylated arsenicals in rat and human cells. *Arch. Toxicol.* 74: 289–299.
- Torloni MR, Betrán AP, Horta BL, Nakamura MU, Atallah AN, Moron AF, et al. 2009. Prepregnancy BMI and the risk of gestational diabetes: a systematic review of the literature with meta-analysis. *Obes Rev* 10:194–203; doi:10.1111/j.1467-789X.2008.00541.x.
- Tsang V, Fry RC, Niculescu MD, Rager JE, Saunders J, Paul DS, et al. 2012. The epigenetic effects of a high prenatal folate intake in male mouse fetuses exposed in utero to arsenic. *Toxicol. Appl. Pharmacol.* 264:439–450; doi:10.1016/j.taap.2012.08.022.
- Tseng CH. 2009. A review on environmental factors regulating arsenic methylation in humans. *Toxicol. Appl. Pharmacol.* 235: 338–350.
- Vahter M. 2002. Mechanisms of arsenic biotransformation. *Toxicology* 181: 211–217.
- Valenzuela OL, Borja-Aburto VH, Cruz-Gonzalez MB, Garcia-Montalvo EA, Calderon-Aranda ES, Del Razo LM. 2005. Urinary trivalent methylated arsenic species in a population chronically exposed to inorganic arsenic. *Environ. Health Perspect.* 113:250–254; doi:10.1289/ehp.7519.
- Vall O, Gómez-Culebras M, Garcia-Algar O, Joya X, Velez D, Rodriguez-Carrasco E, et al. 2012. Assessment of prenatal exposure to arsenic in Tenerife Island. *PLoS ONE* 7:e50463; doi:10.1371/journal.pone.0050463.
- Victora CG, Barros FC, Horta BL, Martorell R. 2001. Short-term benefits of catch-up growth for small-for-gestational-age infants. *Int J Epidemiol* 30: 1325–1330.
- Vincent HK, Taylor AG. 2006. Biomarkers and potential mechanisms of obesity-induced oxidant stress in humans. *International Journal of Obesity* 30:400–418; doi:10.1038/sj.ijo.0803177.

- Watanabe T. 2013. Metabolism of arsenic and its toxicological relevance. *Arch. Toxicol.* 87:969–979; doi:10.1007/s00204-012-0904-5.
- Wickett JC, Vernon PA, Lee DH. 2000. Relationships between factors of intelligence and brain volume. *Personality and Individual Differences* 29:1095–1122; doi:10.1016/S0191-8869(99)00258-5.
- Wilhelm-Benartzi CS, Houseman EA, Maccani MA, Poage GM, Koestler DC, Langevin SM, et al. 2012. In utero exposures, infant growth, and DNA methylation of repetitive elements and developmentally related genes in human placenta. *Environ. Health Perspect.* 120:296–302; doi:10.1289/ehp.1103927.
- Wu G, Bazer FW, Cudd TA, Meininger CJ, Spencer TE. 2004. Maternal nutrition and fetal development. *J. Nutr.* 134: 2169–2172.
- Xu W-X, Liu Y, Liu S-Z, Zhang Y, Qiao G-F, Yan J. 2014. Arsenic trioxide exerts a double effect on osteoblast growth in vitro. *Environmental Toxicology and Pharmacology* 38:412–419; doi:10.1016/j.etap.2014.07.010.
- Xu Y, Wang Y, Zheng Q, Li X, Li B, Jin Y, et al. 2008. Association of oxidative stress with arsenic methylation in chronic arsenic-exposed children and adults. *Toxicol. Appl. Pharmacol.* 232:142–149; doi:10.1016/j.taap.2008.06.010.
- Yang C-Y, Chang C-C, Tsai S-S, Chuang H-Y, Ho C-K, Wu T-N. 2003. Arsenic in drinking water and adverse pregnancy outcome in an arseniasis-endemic area in northeastern Taiwan. *Environ. Res.* 91: 29–34.
- Yunus M, Persson LA. 2008. Nutritional status has marginal influence on the metabolism of inorganic arsenic in pregnant Bangladeshi women. *Environ. Health Perspect.* 116:315–21; doi:10.1289/ehp.10639.

Table 1. Selected characteristics of 706 mothers and their infants from a New Hampshire pregnancy cohort for all participants and by maternal pre-pregnancy weight status^a

Variable	All	Normal Weight	Overweight/Obese	<i>P</i> Value ^b
Pre-pregnancy BMI (kg/m ²), mean (SD)	25.3 (4.9)	22.1 (1.6)	29.8 (4.4)	<0.001
Maternal age (years), mean (SD)	31.1 (4.7)	31.4 (4.7)	30.6 (4.7)	0.03
Education level, N (%) ^c				
Less than 11th grade	8 (1.1%)	3 (0.7%)	5 (1.7%)	0.01
High school graduate or equivalent	68 (9.7%)	34 (8.4%)	34 (11.4%)	
Junior college graduate, or some college or technical school	153 (21.8%)	77 (19.1%)	76 (25.4%)	
College graduate	275 (39.1%)	157 (38.9%)	118 (39.5%)	
Post-graduate schooling	199 (28.3%)	133 (32.9%)	66 (22.1%)	
Previous pregnancies, N (%) ^d				
0	281 (39.9%)	172 (42.6%)	109 (36.3%)	0.04
1	271 (38.5%)	151 (37.4%)	120 (40.0%)	
2	96 (13.6%)	58 (14.4%)	38 (12.7%)	
≥3	56 (8.0%)	23 (5.7%)	33 (11.0%)	
Smoking status, N (%)				
Never	624 (88.4%)	357 (87.9%)	267 (89.0%)	0.58
Ever ^e	45 (6.4%)	29 (7.1%)	16 (5.3%)	
Current	37 (5.2%)	20 (4.9%)	17 (5.7%)	
Maternal gestational weight gain ^f (lbs), mean	32.0 (12.1)	33.8 (9.6)	29.5 (14.5)	<0.001
Maternal gestational hyperglycemia, ^g N (%)				
Normal	566 (83.0%)	332 (84.7%)	234 (80.7%)	0.37
IGT	70 (10.3%)	37 (9.4%)	33 (11.4%)	
GDM	46 (6.7%)	23 (5.9%)	23 (7.9%)	
1-Hour blood glucose challenge test results ^g (mg/dL), mean (SD)	113 (28)	112 (27)	114 (29)	0.29

Home water arsenic concentration (µg/L), median (IQR)	0.5 (0.1-2.7)	0.5 (0.1-2.6)	0.5 (0.1-2.8)	0.94
Any seafood in 2 days prior to urine collection, N (%)	112 (15.9%)	69 (17.0%)	43 (14.3%)	0.39
Any rice in 2 days prior to urine collection, N (%)	163 (23.1%)	102 (25.1%)	61 (20.3%)	0.16
Urinary arsenic concentration (µg/L), median (IQR)				
Inorganic Arsenic (iAs)	0.3 (0.1-0.5)	0.2 (0.1-0.5)	0.3 (0.1-0.5)	0.49
Monomethylarsonic acid (MMA)	0.3 (0.1-0.5)	0.3 (0.1-0.5)	0.3 (0.1-0.6)	0.41
Dimethylarsinic acid (DMA)	2.8 (1.4-4.9)	2.5 (1.3-4.8)	2.9 (1.4-5.1)	0.27
Total Arsenic (iAs+MMA+DMA)	3.4 (1.7-6.0)	3.3 (1.6-5.7)	3.6 (1.8-6.1)	0.34
Primary Methylation Index (PMI) (MMA/iAs)	1.0 (0.5-1.8)	1.0 (0.5-1.8)	1.0 (0.5-1.8)	0.84
Secondary Methylation Index (SMI)	10.4 (7.6-18.3)	10.3 (7.6-17.7)	10.6 (7.5-18.6)	0.68
Infant delivery mode, N (%)				
Vaginal	472 (66.9%)	291 (71.7%)	181 (60.3%)	<0.01
Caesarian Section	234 (33.1%)	115 (28.3%)	119 (39.7%)	
Infant birth outcomes				
Gestational age (weeks), mean (SD)	39.5 (1.6)	39.6 (1.5)	39.3 (1.7)	0.02
Head circumference (cm), mean (SD)	34.7 (1.7)	34.5 (1.8)	34.9 (1.7)	<0.01
Birth length (cm), mean (SD)	50.9 (2.7)	50.9 (2.6)	50.9 (2.8)	0.92
Birth weight (g), mean (SD)	3462 (516.8)	3435 (488.6)	3499 (551.5)	0.11
Ponderal Index (kg/m ³), mean (SD)	26.3 (3.4)	26.1 (3.3)	26.6 (3.6)	0.07
Birth weight status, N (%) ^b				
Small for gestational age	37 (5.2%)	24 (5.9%)	13 (4.3%)	0.02
Average for gestational age	600 (85.0%)	353 (87.0%)	247 (82.3%)	
Large for gestational age	69 (9.8%)	29 (7.1%)	40 (13.3%)	

^aNormal weight defined as pre-pregnancy BMI 18.5-24.9 kg/m² and overweight/obese defined as pre-pregnancy BMI ≥ 25kg/m²; 25 underweight mothers (BMI < 18.5 kg/m²) excluded from analysis

^bCalculated from an unpaired, 2-tailed *t* test with an equal variance assumption to analyze the difference in means, a χ^2 test to analyze the difference in proportions, or a Wilcoxon-Mann-Whitney test to analyze the difference in medians by maternal weight status

^cEducation data missing for 3 participants

^dParity data missing for 2 participants

^eParticipants who reported ever smoking but did not report smoking during pregnancy or had missing data for smoking during pregnancy were classified as ever smokers.

^fData for gestational weight gain available for a subset of 639 women.

^gMaternal hyperglycemia measured as part of usual prenatal care. Results for 1-hour blood glucose screening and IGT status available for a subset of 680 women; data on GDM status available for a subset of 682 women.

^hAge- and sex-adjusted birth weight percentiles were used to classify infants' size-for-gestational age, which included small ($\leq 10^{\text{th}}$ percentile) (SGA), average (11-89th percentile) (AGA) and large-for-gestational age ($\geq 90^{\text{th}}$ percentile) (LGA) (Fenton et al. 2013).

Table 2. Adjusted parameter estimates (95% confidence interval) for the change in birth outcomes with each unit increase in natural log transformed maternal 2nd trimester urinary arsenic for all participants and by maternal pre-pregnancy weight status (N=706)^{a,b}

Outcome/ Maternal Urinary Arsenic Variable	Overall (N=706)	Normal Weight (N=406)	Overweight/Obese (N=300)	P for interaction^c
Gestational age (weeks)^d				
Ln(iAs)	0.04 (-0.06, 0.15)	0.10 (-0.03, 0.23)	-0.04 (-0.20, 0.12)	0.18
Ln(MMA)	0.05 (-0.04, 0.14)	0.08 (-0.03, 0.20)	0.00 (-0.13, 0.14)	0.38
Ln(DMA)	0.09 (-0.03, 0.21)	0.11 (-0.05, 0.27)	0.07 (-0.11, 0.26)	0.78
Ln(tAs)	0.10 (-0.03, 0.22)	0.12 (-0.05, 0.28)	0.07 (-0.11, 0.26)	0.74
Head circumference (cm)^e				
Ln(iAs)	-0.08 (-0.19, 0.02)	-0.12 (-0.26, 0.02)	-0.04 (-0.21, 0.13)	0.51
Ln(MMA)	-0.10 (-0.19, -0.01)*	-0.08 (-0.19, 0.04)	-0.14 (-0.28, 0.00)*	0.49
Ln(DMA)	-0.12 (-0.25, 0.00)	-0.13 (-0.29, 0.04)	-0.11 (-0.30, 0.08)	0.90
Ln(tAs)	-0.14 (-0.27, -0.01)*	-0.15 (-0.32, 0.02)	-0.12 (-0.31, 0.07)	0.83
Birth length (cm)^e				
Ln(iAs)	0.06 (-0.10, 0.22)	0.01 (-0.20, 0.22)	0.13 (-0.12, 0.39)	0.46
Ln(MMA)	0.10 (-0.04, 0.24)	0.02 (-0.16, 0.20)	0.21 (0.00, 0.43)	0.18
Ln(DMA)	0.22 (0.03, 0.41)*	0.09 (-0.16, 0.35)	0.38 (0.09, 0.67)*	0.14
Ln(tAs)	0.21 (0.01, 0.40)*	0.07 (-0.19, 0.33)	0.37 (0.08, 0.67)*	0.12
Birth weight (g)^e				
Ln(iAs)	-9.7 (-38.4, 19.0)	-0.5 (-37.9, 37.0)	-22.9 (-67.6, 21.8)	0.45
Ln(MMA)	3.9 (-20.5, 28.3)	19.4 (-12.4, 51.3)	-18.0 (-55.8, 19.8)	0.14
Ln(DMA)	4.1 (-29.5, 37.8)	24.0 (-20.5, 68.4)	-21.9 (-72.8, 29.0)	0.18
Ln(tAs)	-1.3 (-35.8, 33.2)	15.8 (-30.1, 61.8)	-23.1 (-74.9, 28.7)	0.27

Ponderal index (kg/m ³) ^e				
Ln(iAs)	-0.17 (-0.39, 0.05)	-0.01 (-0.30, 0.28)	-0.40 (-0.75, -0.06)*	0.08
Ln(MMA)	-0.13 (-0.32, 0.06)	0.11 (-0.13, 0.35)	-0.47 (-0.76, -0.19)*	<0.01
Ln(DMA)	-0.32 (-0.58, -0.06)*	0.04 (-0.30, 0.38)	-0.79 (-1.18, -0.40)**	<0.01
Ln(tAs)	-0.34 (-0.60, -0.07)*	0.02 (-0.33, 0.37)	-0.79 (-1.18, -0.40)**	<0.01

^aNormal weight defined as pre-pregnancy BMI 18.5 -24.9 kg/m² and overweight/obese defined as pre-pregnancy BMI ≥ 25 kg/m²; 25 underweight mothers (BMI < 18.5 kg/m²) excluded from analysis.

^bResults from a series of regression models fitting each birth outcome on each arsenic exposure measure (natural log transformed inorganic arsenic (iAs), monomethylarsonic acid (MMA), dimethylarsinic acid (DMA), total arsenic (iAs+MMA+DMA) (tAs)); models are adjusted as noted.

^cP-value from a two-sided Wald's t-test testing the significance of a multiplicative term of maternal pre-pregnancy weight status (categorical) and each arsenic exposure in the regression model.

^dModel adjusted for maternal age (continuous), education level (ordinal with the five categories listed in Table 1 assigned values of 1–5, respectively), parity (continuous), and infant sex; models for overall sample additionally adjusted for maternal pre-pregnancy weight status (categorical).

^eModel adjusted for variables described in footnote d as well as gestational age (continuous)

* $P < 0.05$; ** $P < 0.001$

Table 3. Adjusted parameter estimates (95% confidence interval) for the change in birth outcomes with each unit increase in natural log transformed maternal 2nd trimester urinary arsenic by maternal pre-pregnancy weight status^a and infant sex strata.^b

Maternal pre-pregnancy weight status	Infant Sex	Arsenic Exposure	β (95% Confidence Interval)		
			Birth Length (cm)	Birth Weight (g)	Ponderal index (kg/m ³)
Normal Weight	Female (N=204)	Ln(iAs)	-0.04 (-0.36, 0.27)	-7.1 (-59.3, 45.2)	0.00 (-0.40, 0.40)
		Ln(MMA)	0.02 (-0.24, 0.29)	14.9 (-29.2, 58.9)	0.05 (-0.29, 0.39)
		Ln(DMA)	0.03 (-0.36, 0.42)	34.5 (-29.2, 98.3)	0.17 (-0.32, 0.66)
		Ln(tAs)	0.00 (-0.40, 0.40)	23.5 (-42.4, 89.4)	0.13 (-0.38, 0.63)
	Male (N=202)	Ln(iAs)	0.04 (-0.24, 0.32)	-0.7 (-50.9, 49.4)	-0.04 (-0.45, 0.36)
		Ln(MMA)	0.04 (-0.20, 0.27)	20.5 (-22.0, 63.0)	0.12 (-0.23, 0.46)
		Ln(DMA)	0.17 (-0.16, 0.49)	11.6 (-45.8, 69.0)	-0.12 (-0.58, 0.34)
		Ln(tAs)	0.13 (-0.20, 0.47)	3.7 (-56.1, 63.4)	-0.14 (-0.62, 0.34)
Overweight/Obese	Female (N=153)	Ln(iAs)	0.00 (-0.32, 0.32)	-58.6 (-115.2, -1.9)	-0.50 (-0.97, -0.03)
		Ln(MMA)	-0.07 (-0.38, 0.25)	-85.0 (-139.2, -30.8)**	-0.58 (-1.04, -0.12)
		Ln(DMA)	0.00 (-0.41, 0.40)	-90.7 (-160.6, -20.8)*	-0.69 (-1.27, -0.10)
		Ln(tAs)	0.01 (-0.40, 0.41)	-90.7 (-161.0, -20.5)*	-0.73 (-1.32, -0.14)
	Male (N=147)	Ln(iAs)	0.34 (-0.09, 0.77)	16.9 (-65.2, 98.9)	-0.41 (-0.96, 0.15)
		Ln(MMA)	0.45 (0.14, 0.76)**	28.6 (-31.2, 88.4)	-0.50 (-0.90, -0.09)
		Ln(DMA)	0.75 (0.32, 1.17)**	39.2 (-44.5, 122.9)	-0.95 (-1.50, -0.39)
		Ln(tAs)	0.74 (0.30, 1.19)**	38.7 (-47.7, 125.0)	-0.94 (-1.51, -0.36)
<i>P</i> for 3-way interaction between infant sex, maternal weight status and Ln(tAs) ^c			0.13	0.03	0.89

^aNormal weight defined as pre-pregnancy BMI 18.5 -24.9 kg/m² and overweight/obese defined as pre-pregnancy BMI ≥ 25 kg/m²; 25 underweight mothers (BMI < 18.5 kg/m²) excluded from analysis.

^bResults from a series of regression models fitting each birth outcome on each arsenic exposure measure (natural log transformed inorganic arsenic (iAs), monomethylarsonic acid (MMA), dimethylarsinic acid (DMA), total arsenic (iAs+MMA+DMA) (tAs)). All

models adjusted for maternal age (continuous), education level (ordinal with the five categories listed in Table 1 assigned values of 1–5, respectively), parity (continuous), and gestational age (continuous). Models adjusted for maternal age (continuous), education level (ordinal with the five categories listed in Table 1 assigned values of 1–5, respectively), parity (continuous), gestational age (continuous) and infant sex.

^cTo test for a 3-way interaction between maternal weight status, infant sex, and Ln(tAs), regressions were fit as described in footnote b with the addition of six 2-way multiplicative interaction terms between maternal weight status, infant sex, and arsenic exposure, and one 3-way multiplicative interaction term between all 3 variables; *P*-value from a two-sided Wald *t*-test on the coefficient for the 3-way multiplicative term. Results for test of interaction with Ln(MMA) and Ln(DMA) are similar (data not shown)

P*<0.05; *P*<0.01